

Low scale $B - L$ extension of the Standard Model at the LHC

Shaaban Khalil

Center for Theoretical Physics, British University in Egypt, Cairo, 11837, Egypt.

Department of Mathematics, Faculty of Science, Ain Shams University, Cairo, 11566, Egypt.

Abstract

The fact that neutrinos are massive indicates that the Standard Model (SM) requires extension. We propose a low energy ($\lesssim TeV$) $B - L$ extension of the SM, which is based on the gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$. We show that this model provides a natural explanation for the presence of three right-handed neutrinos in addition to an extra gauge boson and a new scalar Higgs. Therefore, it can lead to very interesting phenomenological implications different from the SM results which can be tested at the LHC. Also we analyze the muon anomalous magnetic moment in this class of models. We show that one-loop with exchange Z' may give dominant new contribution $\sim few \times 10^{-11}$.

1 Introduction

There are at present two pieces of evidence which hint at physics beyond the SM: (i) The solid evidence for neutrino oscillations, pointing towards non-vanishing neutrino masses. In the SM, neutrinos are massless due to the absence of right-handed neutrinos and the exact $B - L$ conservation. (ii) The strength of CP violation in the SM is not sufficient to generate the observed baryon asymmetry in the universe. The neutrino mass puzzle and the baryon asymmetry problem can be readily solved by introducing right-handed neutrinos. Right-handed neutrinos lead to the see-saw mechanism. This mechanism explains, in an elegant way, why neutrinos are much lighter than the other elementary fermions. Also, the new complex phases in the leptonic sector can generate lepton asymmetry, which is converted to baryon asymmetry, through the decay of the right-handed neutrinos.

The tremendous success of gauge symmetry in describing the SM makes us believe that any extension of this model should be through the extension of its gauge symmetry. The minimal type of this extension is based on the gauge group $G_{B-L} \equiv SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$. In fact, the SM is characterized by possessing, at the renormalizable

level, a global $U(1)_{B-L}$ symmetry. If this symmetry is gauged, the existence of three SM singlet fermions (we call them right handed neutrinos) are predicted by the anomaly cancellation conditions, as crucial ingredients for the consistency of the model. In addition, the model also contains an extra gauge boson corresponding to $B - L$ gauge symmetry and an extra SM singlet scalar (heavy Higgs). This may change significantly the SM phenomenology and lead to interesting signatures at the LHC.

In fact, it is very difficult to imagine that the new physics beyond the SM that nature adopted for generating neutrino masses will not be manifested in any other low energy process. Indeed, this is the case when three right-handed neutrinos are randomly added to the SM spectrum. However, with this minimal extension of the SM gauge group, one finds that the neutrino masses are strongly related to some other low energy processes. It is worth mentioning that the extra gauge and Higgs bosons contained in this class of models are playing important role in establishing this relation between the mechanism of generating the neutrino masses and low energy physics consequences.

In this letter we reappraise the low scale scenario of $B - L$ extension of the SM. We will show that this class of model can account for the experimental results of the light neutrino masses and their large mixing. The TeV scale $B - L$ symmetry breaking and see-saw mechanism have not been considered in much details in the literature. There were some attempts in the past for analyzing the $B - L$ extension of the SM [1]. However, these attempts were focused on the neutrino masses, based on old experimental results, in minimal or non-minimal extensions of the SM that include $B - L$ symmetry. Here, we perform a complete analysis for the low scale ($\lesssim 1\text{TeV}$) $B - L$ minimal extension of the SM, where the effects of the new spectrum in different sectors are simultaneously taken into account.

The paper is organized as follows. In section 2 we discuss the $B - L$ extension of the SM with especial emphasis for the simultaneous breaking of $U(1)_{B-L}$ and $SU(2)_L \times U(1)_Y$. The analysis of the Higgs bosons in this model is given in section 3. Section 4 is devoted for the neutrino masses and mixing in this low scale $B - L$ extension of the SM. In section 5 we discuss the new gauge boson Z' in this model and its implication on the muon anomalous magnetic moment. Finally, we give our conclusions in section 6.

2 $B - L$ extension of the SM

In this section we discuss the particle content and the spontaneous symmetry breaking in the minimal extension of the SM based on the gauge group $G_{B-L} \equiv SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$. The invariance of the lagrangian under this gauge symmetry implies the existence of a new gauge boson (beyond the SM ones) C_μ . Also in order to ensure that $U(1)_{B-L}$ is anomaly free, three SM singlet fermions must be introduced. These singlet

fermions are usually called right-handed neutrinos and denoted by ν_{R_i} . In this respect, the global $(B - L)$ symmetry in the SM is gauged and the scale of breaking of this symmetry provides a natural identification to what is called seesaw scale.

The Lagrangian of the leptonic sector in the minimal extension of the SM G_{B-L} is given by

$$\begin{aligned}\mathcal{L}_{B-L} = & -\frac{1}{4}C_{\mu\nu}C^{\mu\nu} + i\bar{l}D_\mu\gamma^\mu l + i\bar{e}_R D_\mu\gamma^\mu e_R + i\bar{\nu}_R D_\mu\gamma^\mu \nu_R + (D^\mu\phi)(D_\mu\phi) \\ & + (D^\mu\chi)(D_\mu\chi) - V(\phi, \chi) - \left(\lambda_e\bar{l}\phi e_R + \lambda_\nu\bar{l}\tilde{\phi}\nu_R + \frac{1}{2}\lambda_{\nu_R}\bar{\nu}^c_R\chi\nu_R + h.c.\right),\end{aligned}\quad (1)$$

where $C_{\mu\nu} = \partial_\mu C_\nu - \partial_\nu C_\mu$ is the field strength of the $U(1)_{B-L}$. The covariant derivative D_μ is generalized by adding the term $ig''Y_{B-L}C_\mu$, where g'' is the $U(1)_{B-L}$ gauge coupling constant and Y_{B-L} is the $B - L$ quantum numbers of involved particles. The Y_{B-L} for leptons and Higgs are given by: $Y_{B-L}(l) = -1$, $Y_{B-L}(e_R) = -1$, $Y_{B-L}(\nu_R) = -1$, $Y_{B-L}(\phi) = 0$ and $Y_{B-L}(\chi) = 2$. In Eq.(1), λ_e , λ_ν and λ_{ν_R} refer to 3×3 Yakawa matrices.

The Higgs sector of this model should contain one $SU(2)_L$ singlet complex scalar field $\chi \equiv (1, 1, 0, 2)$ that can spontaneously break the $U(1)_{B-L}$ symmetry and one $SU(2)_L$ doublet $\phi \equiv (1, 2, 1, 0)$ to break the $SU(2)_L \times U(1)_Y$ symmetry down to $U(1)_{em}$. The non-vanishing vacuum expectation value (vev) of χ : $|\langle\chi\rangle| = v'/\sqrt{2}$ is assumed to be larger than the vev of the Higgs field ϕ : $|\langle\phi^0\rangle| = v/\sqrt{2}$. It is remarkable that the scale of $U(1)_{B-L}$ symmetry breaking, v' , is not fixed. However, as we will see, the mass of the $U(1)_{B-L}$ gauge boson C_μ is given in terms of v' . Therefore it can be bounded from below by the experimental search for extra neutral gauge boson.

In order to analyze the $B - L$ and electroweak symmetry breaking, we consider the most general Higgs potential invariant under these symmetries, which is given by

$$\begin{aligned}V(\phi, \chi) = & m_1^2\phi^\dagger\phi + m_2^2\chi^\dagger\chi + \lambda_1(\phi^\dagger\phi)^2 + \lambda_2(\chi^\dagger\chi)^2 \\ & + \lambda_3(\chi^\dagger\chi)(\phi^\dagger\phi),\end{aligned}\quad (2)$$

where $\lambda_3 > -2\sqrt{\lambda_1\lambda_2}$ and $\lambda_1, \lambda_2 \geq 0$, so that the potential is bounded from below. This is the stability condition of the potential. Furthermore, in order to avoid that $\langle\phi\rangle = \langle\chi\rangle = 0$ be a local minimum, we have to require that $\lambda_3^2 < 4\lambda_1\lambda_2$. As in the usual Higgs mechanism of the SM, the vevs v and v' can not be emerged unless negative squared masses, $m_1^2 < 0$ and $m_2^2 < 0$, are assumed. In this case, the non-zero minimum is given by

$$v^2 = \frac{4\lambda_2m_1^2 - 2\lambda_3m_2^2}{\lambda_3^2 - 4\lambda_1\lambda_2}, \quad v'^2 = \frac{-2(m_1^2 + \lambda_1v^2)}{\lambda_3}.\quad (3)$$

As can be seen from the above expressions, the vevs v and v' can not be emerged unless negative squared masses, $m_1^2 < 0$ and $m_2^2 < 0$, are assumed, as in the usual Higgs mechanism of the SM. It is also interesting to note that with $\lambda_3 > 0$, it is possible to




Figure 1: The $B - L$ and electroweak symmetry breaking minima with $m_1^2 < 0$, $m_2^2 < 0$ and $\lambda_3 \sim -0.5\sqrt{\lambda_1\lambda_2}$.

generate non-vanishing vev v' , *i.e.*, the $B - L$ symmetry is broken, while the vev $v = 0$. This type of minimum corresponds to the scenario of two stages symmetry breaking at different scale, with $v' \gg v$. In our analysis, we are interested in the case of low scale v' , which might be of order the electroweak. Therefore, we will focus on the following region of mixing coupling λ_3 : $0 > \lambda_3 > -2\sqrt{\lambda_1\lambda_2}$. In Fig1. we present the vevs of the potential $V(\phi, \chi)$ in case of $m_{1,2}^2 < 0$, $\lambda_3 \simeq -0.5\sqrt{\lambda_1\lambda_2}$ and $\lambda_{1,2} \sim \mathcal{O}(1)$

After the $B - L$ gauge symmetry breaking, the gauge field C_μ (will be called Z' in the rest of the paper) acquires the following mass:

$$M_{Z'}^2 = 4g''^2 v'^2. \quad (4)$$

The high energy experimental searches for an extra neutral gauge boson impose lower bounds on this mass. The CDF limit [2] leads to $M_{Z'} \gtrsim O(600 - 800)$ GeV. However, since LEP II was e^+e^- collider, it constrain strongly the extra-gauge boson that coupled significantly with electrons. Therefore, LEP II provides the most stringent constraint on $B - L$ gauge boson and implies that [8]*)

$$M_{Z'}/g'' > 6 \text{ TeV}. \quad (5)$$

This implies that $v' \gtrsim O(\text{TeV})$. Moreover, if the coupling $g'' < O(1)$, one can still obtain $m_{Z'} \gtrsim O(600)$ GeV.

*)We would like to thank A. Dobrescu for drawing our attention to this limit.

3 Higgs in $B - L$ extension of the SM

Now we turn to the Higgs sector in this class of models. As mentioned, the Higgs scalar fields in G_{B-L} consists of one complex $SU(2)_L$ doublet and one complex scalar single, *i.e.*, six scalar degrees of freedom. When the $B - L$ and electroweak symmetries are broken, four of them are eaten by Z' , Z^0 and $W^\pm$ bosons and two scalar bosons (ϕ, χ) remain as physical degrees of freedom. From the mass terms in the scalar potential, one finds the following mass matrix for ϕ and χ :

$$\frac{1}{2}M^2(\phi, \chi) = \begin{pmatrix} \lambda_1 v^2 & \frac{\lambda_3}{2} v v' \\ \frac{\lambda_3}{2} v v' & \lambda_2 v'^2 \end{pmatrix}. \quad (6)$$

Therefore, the mass eigenstates fields H and H' are given as

$$\begin{pmatrix} H \\ H' \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \phi \\ \chi \end{pmatrix}, \quad (7)$$

where the mixing angle θ is defined by

$$\tan 2\theta = \frac{|\lambda_3| v v'}{\lambda_1 v^2 - \lambda_2 v'^2}. \quad (8)$$

The masses of H and H' are given by

$$m_{H,H'}^2 = \lambda_1 v^2 + \lambda_2 v'^2 \mp \sqrt{(\lambda_1 v^2 - \lambda_2 v'^2)^2 + \lambda_3^2 v^2 v'^2}. \quad (9)$$

We call H and H' light and heavy Higgs bosons respectively. From these expressions, it is clear that λ_3 is the measuring of the mixing between the SM Higgs and the $B - L$ extra Higgs. For instance, with $\lambda_3 = 0$, there is no mixing and the Higgs masses is given by $m_\phi = \sqrt{2\lambda_1}v$, as in the SM, and $m_\chi = \sqrt{2\lambda_2}v'$. While for $\lambda_3 \neq 0$, one finds that the light Higgs mass becomes smaller than the SM prediction.

Due to the mixing between the two Higgs bosons, the usual SM couplings among the SM-like Higgs H and the SM fermions and gauge bosons are modified as follows:

$$\begin{aligned} g_{Hff} &= i \frac{m_f}{v} \cos \theta, \\ g_{HVV} &= -2i \frac{M_V^2}{v} \cos \theta, \\ g_{HHVV} &= -2i \frac{M_V^2}{v^2} \cos^2 \theta. \end{aligned} \quad (10)$$

Here, we follow the conventions of Ref. [3] for the Feynman rules of the vertices. In addition, there are new couplings among the extra Higgs, H' , and the SM particles :

$$g_{H'ff} = i \frac{m_f}{v} \sin \theta,$$

$$\begin{aligned}
g_{H'VV} &= -2i \frac{M_V^2}{v} \sin \theta, \\
g_{HH'VV} &= -2i \frac{M_V^2}{v^2} \cos \theta \sin \theta.
\end{aligned} \tag{11}$$

We have adopted the first order approximation of the mixing angle θ . Therefore, terms of order $\sin^n \theta$, $n \geq 2$ have been neglected. Furthermore, the right-handed neutrino and the $B - L$ gauge boson Z' are now coupled with both H and H' as follows:

$$\begin{aligned}
g_{H\nu_R\nu_R} &= -i \frac{m_{\nu_R}}{v'} \sin \theta, & g_{H'\nu_R\nu_R} &= i \frac{m_{\nu_R}}{v'} \cos \theta, \\
g_{H'Z'Z'} &= -2i \frac{M_{Z'}}{v'} \cos \theta, & g_{HZ'Z'} &= 2i \frac{M_{Z'}}{v'} \sin \theta, \\
g_{H'H'Z'Z'} &= -2i \frac{M_{Z'}^2}{v'^2} \cos^2 \theta & g_{HH'Z'Z'} &= 2i \frac{M_{Z'}^2}{v'^2} \cos \theta \sin \theta.
\end{aligned} \tag{12}$$

Finally we consider the Higgs self interaction vertices. One can easily prove that they are given by

$$\begin{aligned}
g_{H^3} &= 6i \left(\lambda_1 v \cos^3 \theta - \frac{\lambda_3}{2} v' \cos^2 \theta \sin \theta \right), \\
g_{H'^3} &= 6i \left(\lambda_2 v' \cos^3 \theta + \frac{\lambda_3}{2} v \cos^2 \theta \sin \theta \right), \\
g_{H^4} &= 6i \lambda_1 \cos^4 \theta, & g_{H'^4} &= 6i \lambda_2 \cos^4 \theta, \\
g_{HH'^2} &= 2i \left(\frac{\lambda_3}{2} v \cos^3 \theta + \lambda_3 v' \cos^2 \theta \sin \theta - 3\lambda_2 v' \cos^2 \theta \sin \theta \right), \\
g_{H^2H'} &= 2i \left(\frac{\lambda_3}{2} v' \cos^3 \theta - \lambda_3 v \cos^2 \theta \sin \theta + 3\lambda_1 v \cos^2 \theta \sin \theta \right), \\
g_{H^2H'^2} &= i \lambda_3 \cos^4 \theta.
\end{aligned} \tag{13}$$

These new couplings lead to a different Higgs phenomenology from the well known one predicted by the SM. In Ref.[4], it is shown that in this class of models the cross sections of the SM-like Higgs production are reduced by 20% – 30% in the mass range of 120 – 250 GeV compared to the SM results. While, the implications of the $B - L$ extension to the SM do not change the decay branching ratios. Moreover, the extra Higgs has relatively small cross sections, but it is accessible at LHC.

On the other hand, there are now two fine-tuning problems associated to the electroweak and $B - L$ symmetry breaking. These problems are based on the sensitivity of the Higgs boson masses m_H and $m_{H'}$ to quadratic divergences. It is well known that within the SM, the one loop radiative correction to the Higgs leads to

$$\Delta M_H^2 \propto \frac{3\Lambda^2}{8\pi^2 v^2} \left[M_H^2 + 2M_W^2 + M_Z^2 - 4m_t^2 \right]. \tag{14}$$




Figure 2: Feynman diagrams that lead to one-loop quadratic divergences for the light Higgs boson mass in $B - L$ extension of the SM.

Thus, in order to avoid a fine-tuning between this correction and the tree level value of M_H^2 , an upper bound Λ is obtained. For example for $M_H \simeq 115 - 200$ GeV one finds

$$\left| \frac{\Delta M_H^2}{M_H^2} \right| \leq 10 \quad \rightarrow \quad \Lambda_{\text{SM}} \lesssim 2 - 3 \text{ TeV}. \quad (15)$$

In addition, the cancellation of the quadratic divergences may lead to a prediction for the Higgs mass as suggested by Veltman [5]. Indeed with $M_H \sim (320 \text{ GeV})$ the above correction would vanish. However, this conclusion valid only at the one loop level and one has to ensure that it can also be valid at higher orders.

In the $B - L$ extension of the SM, the Feynman diagrams that lead to one-loop quadratic divergences for the light Higgs boson mass are given in Fig. 1, where Higgs boson, gauge boson and fermion are running in the loop. It is worth noting that three new diagrams different from the SM ones are now involved. These diagrams include loops with right-handed neutrino, Z' gauge boson and heavy Higgs boson. In this case, one obtains

$$\begin{aligned} \Delta M_H^2 &\propto \frac{3\Lambda^2}{8\pi^2} \left[\left(2\lambda_1 + \frac{\lambda_3}{3} \right) \cos^4 \theta + \left(\frac{2M_W^2 + M_Z^2}{v^2} \right) \cos^2 \theta \right. \\ &\quad \left. + \frac{M_{Z'}^2}{v'^2} \sin^2 \theta - 4 \left(\frac{m_t^2}{v^2} \cos^2 \theta + \frac{m_{\nu_R}^2}{v'^2} \sin^2 \theta \right) \right]. \end{aligned} \quad (16)$$

Now the condition of no fine tuning (with $m_H \simeq 200 \text{ GeV}$, $v' \simeq 1 \text{ TeV}$ and $\cos^2 \theta \simeq 0.8$) implies that

$$\Lambda_{\text{SM}} \approx \mathcal{O}(10) \text{ TeV}. \quad (17)$$

The exact limit depends on the values of the masses $M_{Z'}$ and M_{ν_R} . In general, in the $B - L$ model, the SM cut off scale becomes higher than the limit obtained in Eq.(19). This new bound is more consistent with the experimental lower bounds usually imposed to suppress higher order operators. Therefore, it is natural to avoid in this class of model, what is called as a little hierarchy problem. Also, now the mass of the SM-like Higgs is not predictable from the cancellation condition of quadratic divergences.

$$\Delta M_{H'}^2 \propto \frac{3\Lambda^2}{8\pi^2} \left[\left(2\lambda_2 + \frac{\lambda_3}{3} \right) \cos^4 \theta + \left(\frac{2M_W^2 + M_Z^2}{v^2} \right) \sin^2 \theta \right]$$

$$+ \frac{M_{Z'}^2}{v'^2} \cos^2 \theta - 4 \left(\frac{m_t^2}{v^2} \sin^2 \theta + \frac{m_{\nu_R}^2}{v'^2} \cos^2 \theta \right) \Bigg]. \quad (18)$$

It is interesting to note that to avoid a fine-tuning between $M_{H'}^2$ and $\Delta M_{H'}^2$, one finds, for $M_{H'} \simeq 1$ TeV, the following bound

$$\left| \frac{\Delta M_{H'}^2}{M_{H'}^2} \right| \leq 10 \quad \rightarrow \quad \Lambda_{B-L} \lesssim 30 \text{ TeV}. \quad (19)$$

Moreover, Veltman cancellation condition for the extra Higgs quadratic divergences leads to (in the limit of very small mixing)

$$M_{H'}^2 \approx 4M_{\nu_R}^2 - M_{Z'}^2. \quad (20)$$

Thus, one may conclude that $M_{H'} \simeq \mathcal{O}(1)$ TeV.

4 Neutrino masses and mixing in low scale $B - L$ extension of the SM

In this section we analyze the neutrino masses and mixing in the low scale gauge $B - L$ extension of the SM. In this class of models, the neutrino masses may be generated through a TeV scale seesaw mechanism.

After $U(1)_{B-L}$ symmetry breaking, the Yukawa interaction term in Eq.(1): $\lambda_{\nu_R} \chi \nu_R \nu_R$ leads, as usual, to right handed neutrino mass: $M_R = \frac{1}{\sqrt{2}} \lambda_{\nu_R} v'$. Also the electroweak symmetry breaking implies Dirac neutrino mass term : $m_D = \frac{1}{\sqrt{2}} \lambda_\nu v$. Therefore, the mass matrix of the left and right-handed neutrino is given by

$$\begin{pmatrix} 0 & m_D \\ m_D & M_R \end{pmatrix}. \quad (21)$$

Since M_R is proportional to v' and m_D is proportional to v *i.e.*, $M_R \gg m_D$, the diagonalization of the mass matrix leads to the following mass for the light and heavy neutrinos respectively:

$$m_{\nu_L} = -m_D M_R^{-1} m_D^T, \quad (22)$$

$$m_{\nu_H} = M_R. \quad (23)$$

Thus, $B - L$ gauge symmetry can explain the presence of three right handed neutrinos and provide a natural framework for the seesaw mechanism. However as mentioned the scale of B-L and hence the mass M_R of ν_R remains arbitrary. It is often assumed a very large scale for $B - L$ symmetry breaking, *i.e.*, $M_R \sim 10^{15}$ GeV in order to explain the

atmospheric and solar neutrino data. It is important to note that such a large scale may be necessary if the Dirac neutrino masses are assumed to be of order $\mathcal{O}(100)$ GeV. However, there is no any low energy evidence that indicates that the Dirac masses should be of that order. On the contrary, if one tries to establish a flavor symmetry between charged and neutral leptons as in quark sector between up and down, one finds that the Dirac neutrino masses must be very small, of order $\mathcal{O}(10^{-4})$ GeV. This implies that M_R of order TeV is quite acceptable.

In our analysis, we adopt the basis where the charged lepton mass matrix and the Majorana mass matrix M_R are both diagonal. Therefore, one can parameterize M_R as follows

$$M_R = M_{R3} \begin{pmatrix} r_1 & 0 & 0 \\ 0 & r_2 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (24)$$

where

$$M_{R3} = |\lambda_{\nu R3}| \frac{v'}{\sqrt{2}} \quad (25)$$

and

$$r_{1,2} = \frac{M_{R1,2}}{M_{R3}} = \left| \frac{\lambda_{\nu R1,2}}{\lambda_{\nu R3}} \right|. \quad (26)$$

As can be seen from Eq.(24) that even if v' is fixed to be of order TeV, the absolute value of M_R is still parameterized by three known parameters. On the other hand, the Dirac mass matrix (if it is real) is given in terms of 9 parameters. Since $U(1)_{B-L}$ can not impose any further constraint to reduce the number of these parameters, the total number of free parameters involved in the light neutrino mass matrix are 12 parameters. The solar and atmospheric neutrino oscillation experiments have provided measurements for the neutrino mass-squared differences and also for the neutrino mixing angles. At the 3σ level, the allowed ranges are [6]:

$$\begin{aligned} \Delta m_{12}^2 &= (7.9 \pm 0.4) \times 10^{-5} \text{eV}^2, \\ |\Delta m_{32}^2| &= (2.4 \pm 0.3) \times 10^{-3} \text{eV}^2, \\ \theta_{12} &= 33.9^\circ \pm 1.6^\circ, \\ \theta_{23} &= 45^\circ, \\ \sin^2 \theta_{13} &\leq 0.048. \end{aligned} \quad (27)$$

Therefore, the number of the experimental inputs are at most six: three neutrino masses (assuming possible ansatzes like hierarchy or degenerate) and three mixing angles (if we assume $\theta_{13} = 0$).

One of the interesting parametrization for the Dirac neutrino mass matrix is given as follows

$$m_D = \sqrt{M_R R} \sqrt{m_\nu^{diag}} U_{MNS}^\dagger, \quad (28)$$

where m_ν^{diag} is the physical neutrino mass matrix and U_{MNS} is the lepton mixing matrix. The matrix R is an arbitrary orthogonal matrix which can be parameterized-in case of real m_D -in terms of three angles. In Eq.(28), the six unknown parameters are now given in terms of three masses in M_R and the three angles in R . In order to fix these angles, one need a flavor symmetry beyond the gauge symmetry which is typically flavor blind. Several types of flavor symmetries have been discussed in the literatures.

5 Extra $B - L$ gauge boson and muon anomalous magnetic moment

As mentioned above, an extra gauge boson corresponding to $B - L$ gauge symmetry is predicted. In fact, there are many models which contain extra gauge bosons [7–10]. These models can be classified into two categories depending on whether or not they arise in a GUT scenario. In some of these models, the Z' and the SM Z are not true mass eigenstates due to mixing. This mixing induces the couplings between the extra Z' boson and the SM fermions. However, there is a stringent experimental limit on the mixing parameter. In our model of $B - L$ extension of the SM, there is no tree-level $Z - Z'$ mixing. Nevertheless, the extra $B - L$ Z' boson and the SM fermions, are coupled through the non-vanishing $B - L$ quantum numbers. In Ref.[4], it was shown that within $B - L$ extension of the SM the branching ratios of $Z' \rightarrow l^+ l^-$ are of order $\sim 20\%$ compared to $\sim 3\%$ of the SM $BR(Z \rightarrow l^+ l^-)$. Hence, searching for Z' is accessible via a clean dilepton signal at LHC.

Here, we consider the impact of the Z' and also H' on the muon anomalous magnetic moment (a_μ). The a_μ has recently been determined with a very high precision. From the the measurement of E821 collaboration at the Brookhaven National Laboratory the average value of a_μ is given by [11]

$$a_\mu^{\text{exp}} = (116592080 \pm 60) \times 10^{-11} \quad (29)$$

This value differs from the SM prediction [12] by 3.4σ :

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (276 \pm 81) \times 10^{-11} \quad (30)$$

In $B - L$ extension of the SM, a_μ can be generated, in addition to the SM contributions, through other one loop diagrams mediated by the new gauge boson Z' and the extra Higgs boson H' as in Fig. 3. However, the H' contribution is proportional to $\lambda_\mu^2 \sin^2 \theta^2$, therefore its contribution is quite small and can be safely neglected. The Z' contribution is given by

$$\Delta a_\mu \simeq \frac{g''^2}{12\pi^2} \frac{m_\mu^2}{M_{Z'}^2} \simeq \frac{m_\mu^2}{48\pi^2 v'^2} \quad (31)$$




Figure 3: Feynman diagrams for muon anomalous magnetic moment with Z' and H' exchanges.

where m_μ is the mass of the muon. For $v' \sim 1$ TeV, one finds that

$$\Delta a_\mu(B-L) \sim 2.1 \times 10^{-11}, \quad (32)$$

which is clearly not enough to account for the deviation between the experimental result and the SM prediction. Therefore, confirming this discrepancy in anomalous magnetic moment a_μ would be useful hint for further new physics beyond this minimal extension of the SM.

6 Conclusions

We have analyzed the minimal extension of the SM, which is based on the TeV scale $B-L$ gauge symmetry. We have shown that this class of models can give a natural explanation for the TeV scale seesaw mechanism and very small neutrino masses. We provided a detail analysis for the simultaneous breaking of the electroweak and $U(1)_{B-L}$ symmetries. We emphasized that the Higgs sector of this model is quite rich with possible significant mixing between the SM-Higgs and the extra-Higgs. This mixing implies interesting implications, which can be tested at the LHC. Also due to the fact the $B-L$ gauge boson Z' has non-vanishing coupling to the SM-leptons, $Z' \rightarrow l^+ l^-$ is a promising channel for the search for Z' at the LHC. Finally, the muon anomalous magnetic moment was considered in this low $B-L$ extension of the SM. Z' exchange one-loop diagram leads to the dominant new contribution. We found that $a_\mu(Z')$ is of order $few \times 10^{-11}$, which can not accommodate the 3.4σ discrepancy between the experimental result and the SM prediction.

Acknowledgment

This work is partially supported by the ICTP through the OEA-project-30. I would like to thank M. Abbas, W. Emam and R. Mohapatra for their useful discussions.

References

- [1] R. N. Mohapatra and R. E. Marshak, Phys. Rev. Lett. **44**, 1316 (1980) [Erratum-
ibid. **44**, 1643 (1980)]; C. Wetterich, Nucl. Phys. B **187**, 343 (1981); W. Buchmuller,
C. Greub and P. Minkowski, Phys. Lett. B **267**, 395 (1991); W. Buchmuller and
T. Yanagida, Phys. Lett. B **302**, 240 (1993).
- [2] F. Abe *et al.* [CDF Collaboration], Phys. Rev. Lett. **79**, 2192 (1997).
- [3] A. Djouadi, “The anatomy of electro-weak symmetry breaking. I: The Higgs boson
in the arXiv:hep-ph/0503172.
- [4] W. Emam and S. Khalil, arXiv:0704.1395 [hep-ph].
- [5] M. J. G. Veltman, Acta Phys. Polon. B **12**, 437 (1981).
- [6] R. N. Mohapatra and A. Y. Smirnov, arXiv:hep-ph/0603118.
- [7] M. Cvetič and S. Godfrey, arXiv:hep-ph/9504216.
- [8] M. Carena, A. Daleo, B. A. Dobrescu and T. M. P. Tait, Phys. Rev. D **70**, 093009
(2004)
- [9] A. Leike, Phys. Rept. **317**, 143 (1999)
- [10] J. L. Hewett and T. G. Rizzo, Phys. Rep. **183**, 139 (1989)
- [11] G. W. Bennett *et al.* [Muon G-2 Collaboration], Phys. Rev. D **73** (2006) 072003.
- [12] K. Hagiwara, A. D. Martin, D. Nomura and T. Teubner, arXiv:hep-ph/0611102.